

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>7</sup> : A01N 59/12, A61L 15/44</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/54593 (43) International Publication Date: 21 September 2000 (21.09.00)</p>
<p>(21) International Application Number: PCT/EP00/02194 (22) International Filing Date: 13 March 2000 (13.03.00) (30) Priority Data: 9905663.2 12 March 1999 (12.03.99) GB (71) Applicant (for all designated States except US): BRISTOL-MYERS SQUIBB COMPANY [US/US]; 345 Park Avenue, New York, NY 10154 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): PARSONS, Dave [GB/GB]; 6 Briar Drive, Heswell, Wirral L60 5RN (GB). JACQUES, Elizabeth [GB/GB]; 9 Cedar Grove, Hoole, Chester CH2 3LQ (GB). BOWLER, Philip [GB/GB]; 8 Woodbridge Close, Appleton, Warrington, Cheshire WA4 5RD (GB). (74) Agent: MAYS, Julie; Bristol-Myers Company Limited, Patent Dept., Swakeleys House, Milton Road, Ickenham, Uxbridge UB10 8NS (GB).</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: IODINE PREPARATION COMPOSITON (57) Abstract  An iodine preparation composition suitable for use on wounds comprising an iodide source, an oxidant and a buffer characterised in that the iodide is held separately from the oxidant until the point of use, and that the buffer is capable of maintaining the pH of the composition at between pH 4.5 and pH 6 so that iodine is generated at a physiologically acceptable dose rate.</p>		